


Supports Decisions

by Comparing Results to Normative Data



- Compares results with 400+ healthy knee individuals (groups same age and gender)
- Dashboards visualize patient's strengths and deficits




- Reports can immediately be saved/printed reducing the efforts of administration
- Objective patient data and automatically generated reports help physiotherapists/physicians to manage patient expectations and to facilitate the communication between care givers

Balance Agility Speed Strength

For more information or a personal demonstration, please contact your local Zimmer Biomet representative.



Copyright ©2017 Zimmer Biomet, Inc. All rights reserved. This material and all the content, artwork, images and names are subject to copyright and other laws for protection of intellectual property. The copying and distribution of this material other than the intended recipient is prohibited without the prior written permission of Zimmer Biomet.



Back-in-Action

Functional Knee Assessment after ACL Surgery



Sensor-based, **Standardised and Validated**^{1,2} Functional Test after ACL Surgery

Assesses patient's complete functional knee capabilities (balance, agility, speed and strength)

Compares results with normative data of healthy knee population with same age and gender

Supports the decision-making on the best rehabilitation strategy and time point to go back to sports/work by looking at objective measurements

Collects and reports data automatically reducing time of administrative and leading to a stronger relationship with patients and other care givers

Increases patient confidence by providing accurate³ results and the possibility to compare results over time

Two Legs Stability



CMJ



One Leg Stability



One Leg CMJ



Plyometric Jump



Parkour



Quick feet



Standard Test Protocol

7 tests to measure balance, agility, speed and strength

Patient is performing the tests by wearing a validated³ sensor or by using a sensor-based balance board

System automatically collects patient-specific data and saves it on a local device or server

Average testing time: 35 minutes



1. Hildebrandt et al; Functional assessments for decision-making regarding return to sports following ACL reconstruction. Part I: development of a new test battery; Knee Surg Sports Traumatol Arthrosc DOI 10.1007/s00167-015-3529-4
2. Herbst et al; Functional assessments for decision-making regarding return to sports following ACL reconstruction. Part II: clinical application of a new test battery; Knee Surg Sports Traumatol Arthrosc
3. Leardini et al. Journal of NeuroEngineering and Rehabilitation 2014, 11:136, <http://www.jneuroengrehab.com/content/11/1/136>